

UNCLASSIFIED

AD NUMBER
AD810490
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; MAR 1967. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Releases Branch, Frederick, MD 21701.
AUTHORITY
Army Biological Defense Research Lab ltr dtd 28 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD

TECHNICAL MANUSCRIPT 356


INFLUENCE OF UNSATURATION
ON FIBRINOLYTIC ACTIVITY OF FATTY ACIDS

Michael J. Surgalla
Earl D. Beesley
Robert R. Brubaker

MARCH 1967

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

DISCLAIMER NOTICE



**THIS DOCUMENT IS BEST
QUALITY AVAILABLE. THE COPY
FURNISHED TO DTIC CONTAINED
A SIGNIFICANT NUMBER OF
PAGES WHICH DO NOT
REPRODUCE LEGIBLY.**

Reproduction of this publication in whole or in part is prohibited except with permission of the Commanding Officer, Fort Detrick, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 356

INFLUENCE OF UNSATURATION ON FIBRINOLYTIC ACTIVITY
OF FATTY ACIDS

Michael J. Surgalla

Earl D. Beesley

Robert R. Brubaker

Medical Bacteriology Division
BIOLOGICAL SCIENCES LABORATORY

Project 1C014501B71A

March 1967

INFLUENCE OF UNSATURATION ON FIBRINOLYTIC ACTIVITY
OF FATTY ACIDSABSTRACT

Long-chain saturated fatty acids are known to accelerate blood clotting and artificial thrombus formation in vitro, and to produce massive thrombosis and death in dogs and mice. We have found that some long-chain fatty acids induce fibrinolytic activity on bovine fibrin films, with some indication that unsaturated acids were most active. Twofold dilutions of the potassium salts of fatty acids were tested in 0.02-ml amounts for ability to lyse unheated bovine fibrin films. An attempt was made to establish the influence of unsaturation and chain length on fibrinolytic activity. The presence of one unsaturated bond appears to make little difference in activity of C16 fatty acids. In C18 and C20 acids, it increases activity roughly tenfold. In C22 and C24 acids it increases activity more than 100-fold. A possible role of fatty acids in regulation of fibrin formation and digestion is suggested.

Long-chain fatty acids have been reported to accelerate blood clotting and artificial thrombus formation in vitro^{1,2} and to cause massive thrombosis and death in dogs.^{3,4} Unsaturated fatty acids are relatively inactive. We have found recently that long-chain fatty acids can induce fibrinolysis on fibrin plates, and in this case it appeared that the unsaturated acids were more active than the saturated. These preliminary findings are confirmed and extended here.

Fibrinolysis was assayed on a modified Astrup plate prepared in the following manner. Fibrin films were prepared by dissolving Armour bovine fibrinogen (2.5 mg/ml) and Parke, Davis bovine thrombin, topical (50 NIH units/ml) in sodium borate buffer pH 7.7.⁵ Fibrinogen solutions were sterilized by filtration and 10-ml volumes were added to 8.5-cm petri dishes. With the plates on a level surface, 0.5 ml of thrombin solution was added dropwise to the fibrinogen while the mixture was gently swirled to assure thorough distribution. Twofold dilutions of potassium salts of fatty acids were placed on fibrin films in 0.02-ml amounts. The lowest concentration that caused complete perforation of the fibrin film after incubation at 37 C for at least 4 hours was taken as the end point. Figure 1 shows the activity of twofold dilutions of sodium myristate. At concentrations of 20, 10, 5, 2.5, and 1.25 mM, 0.02-ml amounts placed on the fibrin film caused its complete perforation.




Figure 1. Activity of Sodium Myristate
on Bovine Fibrin Film.

Saturated fatty acids having 16, 18, 20, 22, and 24 carbons were compared with acids with one unsaturated bond (Table 1).

TABLE 1. INFLUENCE OF UNSATURATION ON FIBRINOLYTIC ACTIVITY

		Active Concentration, mM				
		Test Number				
		1	2	3	4	5
Palmitic	C16:0		5	0.3	0.3	
Palmitoleic	C16:1		0.5	0.3	0.3	
Stearic	C18:0		0.6	1	1	
Oleic	C18:1	0.2	0.06	0.2	0.08	
Elaidic	C18:1		0.06	0.2	0.2	
Arachidic	C20:0		1	5	1	
Eicosenoic	C20:1					0.08
Behenic	C22:0		>10	>10	>10	
Erucic	C22:1		0.03	0.02	0.04	
Lignoceric	C24:0					>10
Nervonic	C24:1					0.08

Palmitic and palmitoleic acids appear to be equally active. Presence of one unsaturated bond in C18 and C20 acids increases activity roughly tenfold. Presence of one unsaturated bond in C22 and C24 acids increases activity well over 100-fold.

The data in Table 2 suggest that although one unsaturated bond increases activity, polyunsaturation may decrease it. The fourth test of linoleic acid illustrates the frustrations encountered occasionally with the assay method.

TABLE 2. INFLUENCE OF POLYUNSATURATION

Fatty Acid		Active Concentration, mM				
		Test Number				
		1	2	3	4	5
Oleic	C18:1	0.2	0.06	0.2	0.08	
Elaidic	C18:1		0.06	0.2	0.2	
Linoleic	C18:2	0.2	0.5	0.3	10	
Linolenic	C18:3		0.1	0.6	0.6	
Eicosenoic	C20:1					0.08
Arachidonic	C20:4			1	1	

Table 3 shows that saturated fatty acids of chain lengths 12 to 20 are fibrinolytic.

TABLE 3. INFLUENCE OF CHAIN LENGTH ON ACTIVITY OF SATURATED ACIDS

Fatty Acid		Active Concentration, mM			
		Test Number			
		1	2	3	4
Lauric	C12	5	10	3	10
Myristic	C14	1	0.6	3	1
Pentadecanoic	C15		1	6	3
Palmitic	C16		5	0.3	0.3
Stearic	C18		0.6	1	1
Nonadecanoic	C19		6	3	1
Arachidic	C20		1	5	1
Behenic	C22		>10	>10	>10

Gans et al.⁶ have suggested that plasminogen activator plays a role in preventing formation of thrombi because increased plasminogen activator occurs simultaneously with hypercoagulability following intravenous administration of endotoxin in dogs. Pig platelet aggregation induced by behenic acid is counteracted by linoleic or linolenic acids in vitro,⁷ and elevated thrombotic tendency (platelet adhesion) in human patients is reduced by ingestion of linolenic acid.⁸ Since fatty acids are non-fibrinolytic on fibrin films heated to destroy plasminogen, activation of plasminogen appears to be involved in fatty acid fibrinolytic activity. It appears, therefore, that attention should be given to the possible involvement of plasminogen activation in regulating early stages of thrombosis.

LITERATURE CITED

1. Poole, J.C.F. 1955. The effect of certain fatty acids on the coagulation of plasma in vitro. *Brit. J. Exp. Pathol.* 36:248-253.
2. Connor, W.E.; Poole, J.C.F. 1961. The effect of fatty acids on the formation of thrombi. *Quart. J. Exp. Physiol.* 46:1-7.
3. Connor, W.E.; Hoak, J.C.; Warner, E.D. 1963. Massive thrombosis produced by fatty acid infusion. *J. Clin. Invest.* 42:860-866.
4. Hoak, J.C. 1963. Structure of thrombi produced by injections of fatty acids. *Brit. J. Exp. Pathol.* 45:44-47.
5. Lewis, J.H.; Ferguson, J.H. 1950. Studies on a proteolytic enzyme system of the blood: I. Inhibition of fibrinolysin. *J. Clin. Invest.* 29:486-490.
6. Gans, H.; Hanson, M.; Krivit, W. 1963. Effect of endotoxin on clotting mechanism: III. On the relationship between hypercoagulability, plasminogen activator activity, and antithrombin activity. *Surgery* 53:792-796.
7. Mahadevan, V.; Singh, H.; Lundberg, W.O. 1966. Effects of saturated and unsaturated fatty acids on blood platelet aggregation in vitro. *Proc. Soc. Exp. Biol. Med.* 121:82-85.
8. Owren, P.A.; Hellem, A.J.; Odegaard, A. 1964. Linolenic acid for the prevention of thrombosis and myocardial infarction. *Lancet* 2:975-979.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Department of the Army Fort Detrick, Frederick, Maryland 21701		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP
3. REPORT TITLE INFLUENCE OF UNSATURATION ON FIBRINOLYTIC ACTIVITY OF FATTY ACIDS		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (Last name, first name, initial) Surgalla, Michael J. Brubaker, Robert R. Beesley, Earl D.		
6. REPORT DATE March 1967	7a. TOTAL NO. OF PAGES 10	7b. NO. OF REFS 8
8a. CONTRACT OR GRANT NO. b. PROJECT NO. 1C014501B71A c. d.	9a. ORIGINATOR'S REPORT NUMBER(S) Technical Manuscript 356	
9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)		
10. AVAILABILITY/LIMITATION NOTICES Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Department of the Army Fort Detrick, Frederick, Maryland 21701
13. ABSTRACT Long-chain saturated fatty acids are known to accelerate blood clotting and artificial thrombus formation in vitro, and to produce massive thrombosis and death in dogs and mice. We have found that some long-chain fatty acids induce fibrinolytic activity on bovine fibrin films, with some indication that unsaturated acids were most active. Twofold dilutions of the potassium salts of fatty acids were tested in 0.02-ml amounts for ability to lyse unheated bovine fibrin films. An attempt was made to establish the influence of unsaturation and chain length on fibrinolytic activity. The presence of one unsaturated bond appears to make little difference in activity of C16 fatty acids. In C18 and C20 acids, it increases activity roughly tenfold. In C22 and C24 acids it increases activity more than 100-fold. A possible role of fatty acids in regulation of fibrin formation and digestion is suggested.		
14. Key Words *Fatty acids *Fibrinolysis Unsaturated Fibrin		

DD FORM 1473

Unclassified

Security Classification

DISTRIBUTION LIST

<u>ADDRESSEE</u>	<u>NUMBER OF COPIES</u>	<u>ADDRESSEE</u>	<u>NUMBER OF COPIES</u>
Technical Director Building 812	1	Chief, Product Development Division Building 469	1
Chief, Plans and Readiness Operations Office Building 812	1	Commanding Officer U.S. Army Pine Bluff Arsenal ATTN: Director, Biological Operations Pine Bluff, Arkansas 71601	1
Director, Biological Sciences Laboratory Building 560	1	Liaison Representative Animal Disease Investigations Building 1301	1
Chief, Physical Science Division Building 568	2	Liaison Officer U.S. Public Health Service Building 1301	12
Chief, Virus and Rickettsia Division Building 539	1	Commanding Officer U.S. Naval Unit Building 125	3
Chief, Medical Investigation Division Building 604	1	U.S. Army Medical Unit ATTN: Library Building 120	1
Chief, Applied Aerobiology Division Building 568	1	Director U.S. Army Munitions Command Operations Research Group Edgewood Arsenal, Maryland 21010	1
Chief, Test Sphere Branch Applied Aerobiology Division Building 568	1	Commanding Officer U.S. Army Edgewood Arsenal ATTN: SMUEA-TSTI (3) Edgewood Arsenal, Maryland 21010	2
Chief, Test Chamber Branch Applied Aerobiology Division Building 1412	1	Commanding General U.S. Army Munitions Command ATTN: AMSMU-SS-CS Dover, New Jersey 07801	1
Chief, Environmental Analysis Office Building 568	1	Commanding General U.S. Army Munitions Command ATTN: AMSMU-RE-RR Dover, New Jersey 07801	1
Chief, Biomathematics Division Building 1422	1	Commanding Officer U.S. Army Dugway Proving Ground ATTN: Tech Plans & Evaluation Tech Info Division Dugway, Utah 84022	1
Chief, Editorial Branch Technical Information Division Building 816	1		
Chief, Technical Library Branch Technical Information Division Building 426	2		
Chief, Technical Releases Branch Technical Information Division Building 426	10		
Chief, Process Development Division Building 469	1		

<u>ADDRESSEE</u>	<u>NUMBER OF COPIES</u>	<u>ADDRESSEE</u>	<u>NUMBER OF COPIES</u>
Commanding General Deseret Test Center ATTN: Technical Library Fort Douglas, Utah 84113	2	Munitions/ TW Defence Research Staff British Embassy 3100 Massachusetts Avenue, N.W. Washington 8, D.C.	3
Commanding General U.S. Army Materiel Command ATTN: AMCRD-RC Washington, D.C. 20315	1	Canadian Liaison Office (CBR) Building 5101 Edgewood Arsenal, Maryland 21010	3
Defense Documentation Center Cameron Station Alexandria, Virginia 22314	20	Australian Embassy ATTN: Lt. Col. Tonkin Australian Army Staff (W) 2001 Connecticut Avenue, N.W. Washington 7, D.C.	2
Detachment 4, RTD (ATCB) Eglin AFB, Florida 32542	1	University of Pennsylvania Institute for Cooperative Research Project Summit 3634 Walnut Street Philadelphia, Pennsylvania 19104	1
APGC(PGBPS-12) Eglin AFB, Florida 32542	1	Institute for Cooperative Research University of Pennsylvania Reference Center P.O. Box 1867 Eglin AFB, Florida 32542	1
USAF EL Lackland AFB, Texas 78236	1	Oak Ridge National Laboratory ATTN: Librarian, Civil Defense Group Oak Ridge, Tennessee 37831	1
Scientific Director Naval Biological Laboratory Naval Supply Center Oakland, California 94614	1	Chief, Medical Bacteriology Division Building 560	10
Commanding Officer and Director U.S. Naval Applied Science Laboratory ATTN: Code 9440 U.S. Naval Base, Brooklyn, N.Y. 11251	1		
Medical Director Naval Ammunition Depot Crane, Indiana 47522	2		
US Army Standardization Group-Canada Office, Senior Standardization Rep. c/o Director of Equipment Policy Canadian Army Headquarters Ottawa 4, Canada	1		